

**CIRCUMFERENCE REDUCTION FOR THE ABDOMEN  
INCLUSION & EXCLUSION CRITERIA  
PLEASE READ CAREFULLY**

**Inclusion Criteria**

- Healthy females or males, between 21 and 60 years of age with a BMI between 18.5 & 29.9
- Those willing to be available for all study visits and complete all procedures.
- Those willing to refrain from a change in diet/drinking/exercise/medication regimen during the course of the study
- If female must be using a medically acceptable form of birth control at least three months prior to enrollment (i.e. condoms with spermicide, IUD, birth control pills)

**Exclusion Criteria**

- Pregnant women, women who are breast feeding, have given birth in the past three (3) months or women who are planning a pregnancy
- Having any electrical implant in the body, such as a pacemaker or an internal defibrillator
- Having a permanent implant in the treatment area, such as metal or injected chemical substance such as silicone
- Having undergone a liposuction surgery or any contouring treatment in the treatment area within three (3) months of study enrollment
- Having received treatment to the treatment area with laser, Radio Frequency or any other device within six (6) months of study enrollment
- Having undergone any other surgeries in the treatment area within nine (9) months
- Having or undergoing any form of treatment for active cancer, or have a history of skin cancer in the treatment area
- Suffering from significant concurrent illness, such as cardiac disorders, diabetes (type I & II), lupus, porphyria, or pertinent neurological disorders
- Having a known anticoagulative or thromboebolic condition or taking anticoagulation medication one (1) week prior to and during treatment course
- History of immunosuppression/ immune deficiency disorders (including HIV or AIDS)
- Suffering from any hormonal imbalance which may affect weight or cellulite such as a thyroid condition that is not under control
- History of lymphatic drainage problems
- Suffering from any significant skin conditions or inflammatory skin conditions in the treatment area
- Having a history of Keloid scarring or abnormal wound healing
- Having a history of epidermal or dermal disorders or being especially prone to bruising
- Any significant change in diet or exercise regimen within one (1) month of enrollment or weight loss and/or gain of  $\geq 10$  lbs within two (2) months of enrollment.
- Use of isotretinoin within 6 months of treatment or during the study